

Original Paper

# The Annual American Men's Internet Survey of Behaviors of Men Who Have Sex With Men in the United States: Protocol and Key Indicators Report 2013

Travis Howard Sanchez<sup>1</sup>, DVM, MPH; R Craig Sineath<sup>1</sup>, MPH; Erin M Kahle<sup>1</sup>, PhD, MPH; Stephen James Tregear<sup>2</sup>, DPhil; Patrick Sean Sullivan<sup>1</sup>, DVM, PhD

<sup>1</sup>Emory University, Atlanta, GA, United States

<sup>2</sup>Manila Consulting Group, McLean, VA, United States

**Corresponding Author:**

Travis Howard Sanchez, DVM, MPH

Emory University

1518 Clifton Road NE

Atlanta, GA, 30322

United States

Phone: 1 404 727 8403

Fax: 1 404 727 8737

Email: [Travis.Sanchez@emory.edu](mailto:Travis.Sanchez@emory.edu)

## Abstract

**Background:** Men who have sex with men (MSM) are disproportionately affected by human immunodeficiency virus (HIV) and there is evidence that this population is participating in increasingly risky sexual behavior. These changes are occurring in the context of new modes of online social interaction—many MSM now report first meeting their sex partners on the Internet. Better monitoring of key behavioral indicators among MSM requires the use of surveillance strategies that capitalize on these new modes of interaction. Therefore, we developed an annual cross-sectional behavioral survey of MSM in the United States, the American Men's Internet Survey (AMIS).

**Objective:** The purpose of this paper was to provide a description of AMIS methods. In addition we report on the first cycle of data collection (December 2013 through May 2014; AMIS-2013) on the same key indicators used for national HIV behavioral surveillance.

**Methods:** AMIS-2013 recruited MSM from a variety of websites using banner advertisements or email blasts. Adult men currently residing in the United States were eligible to participate if they had ever had sex with a man. We examined demographic and recruitment characteristics using multivariable regression modeling ( $P<.05$ ) stratified by the participants' self-reported HIV status.

**Results:** In the AMIS-2013 round, 79,635 persons landed on the study page and 14,899 were eligible, resulting in 10,377 completed surveys from MSM representing every US state. Participants were mainly white, 40 years or older, living in the US South, living in urban areas, and recruited from a general social networking website. Self-reported HIV prevalence was 10.73% ( $n=1113$ ). Compared to HIV-negative/unknown status participants, HIV-positive participants were more likely to have had anal sex without a condom with any male partner in the past 12 months (72.24% versus 61.24%, respectively;  $P<.001$ ) and more likely to have had anal sex without a condom with their last male sex partner who was discordant/unknown HIV status (42.95% versus 13.62%, respectively;  $P<.001$ ). Illicit substance use in the past 12 months was more likely to be reported by HIV-positive participants than HIV-negative/unknown status participants (39.17% versus 26.85%, respectively;  $P<.001$ ). The vast majority of HIV-negative/unknown status participants (84.05%) had been previously HIV tested, but less than half (44.20%) had been tested in the past 12 months. Participants 18-24 years of age were more likely than those 40 years or older to have had anal sex without a condom with a discordant/unknown HIV status partner, were more likely to report substance use, and were less likely to have been HIV tested. Compared to general social networking, those from a geospatial social networking website were more likely to have reported all risk behaviors but were more likely to have been HIV tested.

**Conclusions:** The first round of AMIS generated useful behavioral measures from more than 10,000 MSM Internet users. Preliminary findings identified some subgroups of MSM Internet users that are at potentially higher risk of HIV

acquisition/transmission. AMIS will provide an ongoing data source for examining trends in sexual risk behavior of MSM. This will help to plan and monitor the impact of programs to improve this population's health.

(*JMIR Public Health Surveill* 2015;1(1):e3) doi: [10.2196/publichealth.4314](https://doi.org/10.2196/publichealth.4314)

## KEYWORDS

MSM; gay; homosexual; bisexual; HIV; STD; Internet; survey; surveillance

## Introduction

Men who have sex with men (MSM) continue to be disproportionately affected by human immunodeficiency virus (HIV). In the United States in 2012, more than 30,000 MSM were newly diagnosed with HIV infection, representing 66% of all diagnoses that year [1]. In contrast, gay/bisexual-identifying men account for <2% of the total US population [2]. There is also evidence that risky sexual behavior among MSM has increased in the past decade; data from the Centers for Disease Control and Prevention's National HIV Behavioral Surveillance system (NHBS), that collects data on MSM in major US cities every three years, show a significant increase in the proportion of MSM who reported having anal sex without a condom between their 2005 and 2011 surveys [3]. From 2002-2011, MSM were also the only risk group for whom new HIV diagnoses did not decline [4], and HIV incidence among young MSM is estimated to have increased in recent years [5].

Contemporary to these increased HIV risks are new advances in HIV prevention for MSM. The past 5 years has seen new research proving the efficacy of antiretroviral medication to prevent HIV acquisition (pre-exposure prophylaxis or PrEP) and treatment of HIV positive persons that can reduce transmission [6,7]. Modeling has shown that implementing these biomedical interventions as part of an overall package of HIV prevention services could avert at least one quarter of HIV transmissions among MSM [8]. There are also now more sensitive tests that can detect HIV as early as 1 week after infection and a self-administered rapid HIV test [9,10].

All of these changes are occurring in a new context of social interaction. There are growing numbers of social networking website users and mobile application users [11]. MSM frequently report that they first met their sex partners online and spend considerable time looking for new partners this way [12-15]. This pattern of changing social context for MSM has been capitalized upon by many previous researchers who have successfully conducted entire cross-sectional research studies with MSM online [12-31].

There has also been progress made in large-scale behavioral surveys of MSM designed to monitor key risk behaviors over time. An example of such a system is the Gay Men's Sex Survey that has been conducted with Internet-recruited MSM in the United Kingdom every year since 2001 [32]. The Internet component of the survey now comprises the majority of the more than 10,000 annual survey respondents [33]. The largest ever Internet survey of MSM, the European MSM Internet Survey, was conducted in 2010 and collected data from 180,000 MSM in 38 European countries [34]. This study proved that the

Internet is a viable and useful approach for large-scale behavioral surveillance.

In the United States, there has also been exploration of methods for routine monitoring of HIV-related risk behaviors among MSM. There was a one-time feasibility pilot of the Web-based HIV behavioral surveillance system (WHBS) conducted by the Centers for Disease Control and Prevention as supplement to NHBS [3,15,35]. The primary purpose of WHBS was to conduct behavioral surveillance with a standardized survey to compare to other data collections of MSM and estimate prevalence of risk behaviors among MSM Internet users. The pilot was successful at garnering a large sample of MSM.

There remains a need for establishing a system that can produce data for timely and large-scale monitoring of the behavior trends among MSM. In response to this need, we developed a new annual cross-sectional Internet survey of MSM in the United States, the American Men's Internet Survey (AMIS). The goal of AMIS is to collect surveys from 10,000 MSM each year in the United States in order to generate annual snapshots of relevant behaviors. In this paper, we provide the detailed description of our methods/materials, and report recruitment outcomes and some key indicators from our first round of data collection. To help with comparisons, the key indicators and the analytic approach were designed to mirror those used by NHBS's most recent report on MSM risk behavior [3].

## Methods

### Recruitment and Enrollment

AMIS participants were recruited through convenience sampling from a variety of websites using banner advertisements or email blasts to website members (hereafter referred to generically as "ads"). Ads depicted male models of various races and ethnicities (Figure 1). Men who clicked on the ads were taken directly to the survey website. Two survey platforms were used, depending on the recruitment website. Men recruited through ads posted to a geospatial social networking application were taken to our mobile-optimized survey hosted on a secure server administered by SurveyGizmo (Boulder, Colorado). Men recruited through ads posted elsewhere were taken to our survey hosted on a custom-designed survey website, also hosted on a secure server. Both survey websites used the same study content, used the same security standards, and were compliant with the Health Information Portability and Accountability Act.

The first page that men encountered on the study website contained a brief description of the study. Those who were interested in participating clicked a "begin survey" button that took them to the study's informed consent page which contained standard information regarding the study purpose, procedures, risks, benefits, protections, and investigator contact information.

Those who consented to participate in the study were asked to check a box affirming this decision before continuing. Men who consented were then taken to a page with a brief eligibility screening questionnaire. To be eligible for the survey, participants had to be 18 years of age or older, consider themselves to be male, and report that they had oral or anal sex with a man at least once in the past. As is standard in behavioral

research with MSM, transgender persons were excluded from the study because they are not MSM and recruitment approaches and behavioral risk measures should be specifically designed for this group. Persons who reported being <18 years of age or refused to provide their age were not asked any other screening questions. Persons who reported any gender identity other than male were not asked the sex behavior screening questions.

**Figure 1.** Example Banner Advertisements Used for the American Men's Internet Survey, 2013.



## Survey Administration

MSM who met the eligibility criteria started the online survey immediately. The survey consisted of a core questionnaire administered to all participants, 3 different subset questionnaires to which participants were randomized at the start of the survey, and an additional set of questions that were asked only of participants recruited through geospatial social networking ads. The subset questionnaires were of similar lengths. The intent of the randomization was to reduce overall survey response burden while still generating useful information on some additional behaviors. Participants were blind to this randomization and the randomized subset questions were interspersed with the core questions. The core questions were comprised of the following domains: demographics, sexual behavior, HIV testing history, drug and alcohol use, and HIV prevention services exposure. The randomized question subsets were comprised of the following domains: Subset A—knowledge and use of antiretrovirals for HIV prophylaxis and sexually transmitted disease testing/vaccination; Subset B—disclosure of sexual identity and experiences of stigma; and Subset C—additional details about most recent male sex partner. The participants recruited from the geospatial social networking website received an additional set of questions about a potential mobile HIV prevention app and about acute HIV infection. The core and subset questions were derived from those validated and used by NHBS and used the same timeframes for behaviors [3,36]. The full questionnaire is presented in [Multimedia Appendix 1](#).

To maximize the efficiency of the survey, questions were presented adaptively using a participant's previous responses to determine the path of questioning or auto-filling responses into the text of follow-up questions. On average there were 5 questions per survey webpage. Participants could decline to answer any question. Participants were not asked to correct, verify, or complete responses at the end of their survey. A participant who left their survey would not be able to see any of their previous responses on returning and would have to start the whole survey over again. We did not use cookies to

minimize information on a participant's computer that could potentially identify them as an AMIS participant. We did collect Internet protocol (IP) address to allow us to determine residency and identify potential duplicate respondents.

## Human Subjects Protections

The study was conducted in compliance with federal regulations governing protection of human subjects and was reviewed and approved by our institution's human subjects research review board. No incentive was provided to the participants. Datasets for analyses are stored on secure data servers with access only granted to study staff. The study data are protected under a federal certificate of confidentiality that prevents legal action to force data release.

## Measures and Analyses

Recruitment outcomes for the study are reported as screening, eligibility, unduplicated responses, survey success and reporting sex with a man in the past 12 months. Screening was defined as those who started the screening questionnaire. Overall survey eligibility and individual criteria for ineligibility are presented and were based on survey responses for age, gender and sexual behavior. US residency was determined by either a response of a valid US ZIP code of residence or, for those with no valid ZIP code response, an IP address assigned to a location in the US. Unduplicated responses were determined based on the de-duplication algorithm using IP address, response matching and survey success (see [Multimedia Appendix 2](#) for details). If an observation had a missing value for the first question of at least two consecutive sections, their response was considered incomplete and was not included in the final dataset. All other surveys were considered a "success". Sex with a man in the past 12 months was determined by reporting of one or more partners in response to the question, "In the past 12 months, with how many different men have you had oral or anal sex?"

In addition to standard individual demographic characteristics, we categorized participants based on recruitment source, self-reported HIV status, and geography. The embedded links in the ads were unique and allowed us to determine from which

website participants were recruited. We categorized these based on target audience and purpose: gay social networking ( $n=2$ ), gay general interest ( $n=3$ ), general social networking ( $n=1$ ), and geospatial social networking ( $n=1$ ). We do not provide the names of the websites to preserve operator/client privacy, particularly where a website category has only one operator. Gay social networking websites are those designed for gay or bisexual men to connect with one another, including those attempting to connect for sex. Gay general interest websites are those designed specifically for gay or bisexual men's general interests, such as news stories, public policy advocacy, and travel. The general social networking website is one designed for the general public to connect with others and is not specifically focused on connecting sexual partners. The geospatial social networking website runs on smart cellular telephones and is designed for gay and bisexual men to connect to other men who are near their current location, including those attempting to connect for sex. Self-reported HIV status was determined from responses to questions about having ever had an HIV test, results of the most recent HIV test, and having ever had a positive HIV test. Participants were categorized as HIV-positive, HIV-negative, or unknown status.

We used a combination of county and ZIP code of residence to determine state, US Census-based region, NHBS city residency, and population density. Cities included in the NHBS as of 2011 were as follows: Atlanta, Georgia; Baltimore, Maryland; Boston, Massachusetts; Chicago, Illinois; Denver, Colorado; Houston, Texas; Los Angeles, California; Miami, Florida; Newark, New Jersey; New York City, New York; Philadelphia, Pennsylvania; San Diego, California; San Francisco, California; San Juan, Puerto Rico; Washington, DC; Dallas, Texas; Detroit, Michigan; New Orleans, Louisiana; Nassau-Suffolk, New York; and Seattle, Washington.

The participants who were eligible, unduplicated, successful, and reported male-male sex in the past 12 months were included in analyses of participant characteristics and behavior. Overall chi-square tests were used to identify whether participant characteristics significantly differed between recruitment website types and between question subsets. Following the format used by NHBS in the most recent report of MSM behaviors [3], the prevalence of sex and substance-using behaviors were stratified by self-reported HIV status as either HIV-positive or HIV-negative/unknown. Sexual behaviors were assessed with male partners for either the past 12 months (anal intercourse without a condom with any partner) or for the last partner (anal intercourse without a condom with a discordant or unknown status partner) [3]. HIV serostatus discordance was based on the participant's HIV status and the status of their sex partner. Discordance was defined as either the participant or partner having unknown status or when one was HIV-negative and the other was HIV-positive. Sexually transmitted infection (STI) testing and diagnosis in the past 12 months was only assessed for one-third of randomized participants and included gonorrhea, Chlamydia and syphilis [37]. Illicit substance use in the past 12 months was assessed as the use of any type of illicit substance

by any means of delivery, including injection [37]. Binge alcohol drinking in the past 12 months was assessed as having at least once had 5 or more alcoholic drinks in one sitting [38].

Prevalence of sexual behaviors with male partners, substance use and HIV testing were also presented by race/ethnicity, age group, NHBS city residency, and website recruitment type within the HIV status categories. To determine whether there were significant differences in reported behaviors of different participant subgroups, we conducted multivariable modeling stratified by self-reported HIV status in which each behavior was modeled as the dependent variable and including the following independent variables: race/ethnicity, age group, NHBS city residency, and recruitment website type. We also conducted multivariable logistic regression modeling to determine significant differences in behaviors based on self-reported HIV status while controlling for race/ethnicity, age group, NHBS city residency, and recruitment website type. HIV testing behaviors were only examined among those who did not report that they were HIV-positive and were also presented by participant characteristics. Multivariable logistic regression results are presented as Wald chi-square  $P$  values to denote an independently significant difference in the behavior for each sub-group compared to a referent group. Statistical significance was determined at  $P<.05$ .

## Results

### Recruitment, Enrollment, and Survey Completion

The 2013 data collection round of AMIS (AMIS-2013) ran from December 2013 through May 2014, and resulted in 79,635 persons clicking on the ads and landing on the study's recruitment page (Table 1). Most were from a general social networking website (36,281/79,635, 45.56%) or a geospatial social networking website (27,720/79,635, 34.81%). About a quarter of those who landed on the study's page (18,669/79,635, 23.44%) consented to take part in it. The proportion providing consent varied by recruitment website, with the highest proportion consenting among those recruited from gay general interest websites (36.97%) and the lowest proportion among recruits from the geospatial social networking website (14.18%). Most who were screened were eligible (79.81%). The most common reasons for ineligibility were not being male or reporting not having male-male sex. This was true even of the websites that were specifically marketed to gay persons.

There were 709 (4.76%) surveys determined to likely be from duplicate participants. Among unduplicated surveys, most were considered successful (12,369/14,190, 87.17%). Most successful surveys were among men who reported having sex with another man in the past 12 months (10,377/12,369, 83.90%). The median duration of completion for successful surveys from MSM participants was 14 minutes. AMIS-2013 was managed, implemented, and analyzed by 4 part-time staff (2 faculty, 1 post-doctoral fellow, and 1 program associate). The total cost to implement the survey was approximately 150,000 USD or 15 USD per successful survey.



**Table 1.** Recruitment outcomes with different recruitment website types for the American Men's Internet Survey, United States, 2013.

Recruitment outcomes	Total		Gay social networking (n=2)		General gay interest (n=3)		General social networking (n=1)		Geospatial social networking (n=1)	
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
Clicked ad	79,635		6889		8745		36,281		27,720	
Consented <sup>a</sup>	18,669	(23.44)	1404	(23.38)	3233	(36.97)	10,100	(27.84)	3932	(14.18)
<b>Ineligible<sup>b</sup></b>	3770		223	(15.88)	632	(19.55)	2304	(22.81)	611	(15.54)
Not 18+ years of age <sup>c</sup>	636	(16.87)	54	(24.22)	109	(17.25)	311	(13.50)	162	(26.51)
Not male <sup>c</sup>	2132	(56.55)	175	(78.48)	450	(71.20)	1052	(45.66)	455	(74.47)
Not ever MSM <sup>c</sup>	3628	(96.23)	223	(100)	631	(99.84)	2304	(100.00)	470	(76.92)
Not a resident <sup>c</sup>	1408	(37.35)	119	(53.36)	369	(58.39)	732	(31.77)	188	(30.77)
Eligible <sup>b</sup>	14,899	(79.81)	1181	(84.12)	2601	(80.45)	7796	(77.19)	3321	(84.46)
Unduplicated <sup>d</sup>	14,190	(95.24)	1130	(95.68)	2516	(96.73)	7373	(94.57)	3171	(95.48)
Success <sup>e</sup>	12,369	(87.17)	987	(87.35)	2270	(90.22)	6735	(91.35)	2377	(74.96)
MSM <sup>f</sup> past 12 months <sup>g</sup>	10,377	(83.90)	802	(81.26)	1958	(86.26)	5336	(79.23)	2281	(95.96)

<sup>a</sup> Proportion is of total who clicked ad<sup>b</sup> Proportion is among consented<sup>c</sup> Proportion is among total ineligible<sup>d</sup> Proportion is among eligible. Unduplicated removes participants who were marked as duplicates using IP address and demographic data.<sup>e</sup> Proportion is among unduplicated. Success removes participants who did not pass the survival analysis test for survey completeness.<sup>f</sup> MSM: Men who have sex with men<sup>g</sup> Proportion is among successes

## Participant Characteristics

Of the 10,377 participants in AMIS-2013 who had a successful survey and had male-male sex in the past 12 months, more than three-quarters were white, non-Hispanic (Table 2). Nearly half of the participants were ≥ 40 years of age; others were distributed almost equally between younger age groups. The most common region of residence was the South followed by the West. AMIS-2013 had participants from all US states and at least 100 participants from each of 27 states (Figure 2). There were approximately twice as many participants from urban areas as there were from rural areas, and about one-third of participants lived in NHBS cities. Overall, 1113 (10.73%) participants reported being HIV positive and 9264 (89.27%) reported being HIV negative or having an unknown HIV


serostatus. Most participants were recruited from a general social networking website. The second most common recruitment site was the geospatial social networking website.

There were significant differences in participant characteristics based on where they were recruited (Table 2, all  $P < .001$ ). Most of those differences were observed among participants recruited from the geospatial social networking website, who were less likely be white, less likely be 40 years or older, less likely to live in an NHBS city, more likely to live in the South, more likely to live in urban areas, and more likely to report being HIV positive. There were no significant differences in the characteristics of survey sub-samples that received the 3 different randomized questionnaires (see Multimedia Appendix 3).

**Table 2.** Characteristics of MSM participants in the American Men's Internet Survey by recruitment website type, United States, 2013.

Participant characteristics	Total		Gay social networking (n=2)		General gay interest (n=3)		General social networking (n=1)		Geospatial social networking (n=1)	
	n	(%)	n	(%)	n	(%)	n	(%)	n	(%)
<b>Race/Ethnicity</b>										
Black, non-Hispanic	354	(3.41)	37	(4.61)	48	(2.45)	119	(2.23)	150	(6.58)
Hispanic	1084	(10.45)	66	(8.23)	127	(6.49)	417	(7.81)	474	(20.78)
White, non-Hispanic	8076	(77.83)	645	(80.42)	1638	(83.66)	4351	(81.54)	1442	(63.22)
Other or multiple races	863	(8.32)	54	(6.73)	145	(7.41)	449	(8.41)	215	(9.43)
<b>Age (years)</b>										
18-24	1982	(19.10)	96	(11.97)	246	(12.56)	1067	(20.00)	573	(25.12)
25-29	1515	(14.60)	76	(9.48)	256	(13.07)	693	(12.99)	490	(21.48)
30-39	1918	(18.48)	80	(9.98)	398	(20.33)	881	(16.51)	559	(24.51)
40 or older	4962	(47.82)	550	(68.58)	1058	(54.03)	2695	(50.51)	659	(28.89)
<b>Region</b>										
Midwest	2078	(20.03)	203	(25.31)	347	(17.72)	1147	(21.50)	381	(16.70)
Northeast	2050	(19.76)	176	(21.95)	444	(22.68)	1119	(20.97)	311	(13.63)
South	3558	(34.29)	279	(34.79)	648	(33.09)	1681	(31.50)	950	(41.65)
West	2503	(24.12)	144	(17.96)	518	(26.46)	1382	(25.90)	459	(20.12)
US dependent areas	8	(0.08)	0	(0.00)	1	(0.05)	7	(0.13)	0	(0.00)
<b>NHBS<sup>a</sup> City Resident</b>										
Yes	3268	(31.49)	244	(30.42)	858	(43.82)	1750	(32.80)	416	(18.24)
No	7109	(68.51)	558	(69.58)	1100	(56.18)	3586	(67.20)	1865	(81.76)
<b>Population Density</b>										
Rural	3833	(36.94)	360	(44.89)	626	(31.97)	2129	(39.90)	718	(31.48)
Urban	6544	(63.06)	442	(55.11)	1332	(68.03)	3207	(60.10)	1563	(68.52)
<b>Self-reported HIV Status</b>										
Positive	1113	(10.73)	50	(6.23)	155	(7.92)	439	(8.23)	469	(20.56)
Negative	7657	(73.79)	535	(66.71)	1556	(79.47)	4018	(75.30)	1548	(67.86)
Unknown	1607	(15.49)	217	(27.06)	247	(12.61)	879	(16.47)	264	(11.57)
<b>Total</b>	10,377		802		1958		5336		2281	

<sup>a</sup> NHBS: National HIV Behavioral Surveillance System

**Figure 2.** Number of MSM Participants in the American Men's Internet Survey by State, 2013.

## Sexual Behaviors

Most participants had anal sex without a condom with another man in the past 12 months (Table 3). The proportion who had anal sex without a condom was significantly higher among HIV-positive participants compared to HIV-negative/unknown status participants (72.24% versus 61.24%, respectively;  $P < .001$ ). Compared to HIV-negative/unknown status participants, a larger proportion of HIV-positive participants had anal sex without a condom with their last male sex partner who was discordant/unknown status (13.62% versus 42.95%, respectively;  $P < .001$ ).

Among those who were HIV-positive, Hispanic participants were less likely than white participants to report anal sex without a condom in the past 12 months and black participants were less likely than white participants to report anal sex without a condom with an HIV-negative/unknown status partner (Table 3). Participants 18-24 years of age were more likely to report anal sex without a condom with an HIV-negative/unknown status partner compared to participants  $\geq 40$  years of age. Nearly two-thirds of HIV-positive participants 18-24 years reported anal sex without a condom with a partner who was either HIV-negative or of unknown status. HIV-positive participants who lived in NHBS cities were also more likely than those living elsewhere to report anal sex without a condom in the past 12 months. Compared to HIV-positive participants from a general social networking website, those recruited from a geospatial social networking website were also significantly

more likely to report anal sex without a condom and anal sex without a condom with an HIV-negative or unknown status partner.

Among those who were HIV-negative or unknown status, those 25-39 years of age were significantly more likely to report anal sex without a condom compared to participants  $\geq 40$  years of age (Table 3). Participants 18-24 years of age were more likely to report anal sex without a condom with an HIV-positive or unknown status partner compared to participants  $\geq 40$  years of age. Compared to participants recruited from the general social networking website, those from other websites had significant differences sexual behaviors. Participants from gay social networking websites were less likely to report anal sex without a condom, but those from a geospatial social networking website were more likely to report this behavior. Participants from gay social networking and geospatial social networking websites were more likely to report anal sex without a condom with an HIV-positive/unknown status partner.

Among MSM participants who were HIV-positive, 3.05% (34/1113) also had sex with a woman and 1.17% (13/1113) of those participants reported vaginal sex without a condom in the past 12 months (data not presented in a table). Among those who were HIV-negative or unknown status, 10.29% (953/9264) also had sex with a woman and 6.50% (602/9264) had vaginal sex without a condom in the past 12 months. Both behaviors were significantly more likely among HIV-negative or unknown status participants than among HIV-positive participants (both  $P < .001$ ).

**Table 3.** Sexual Behaviors with Male Partners of MSM Participants in the American Men's Internet Survey, United States, 2013.

Participant characteristics	n in sample	Anal intercourse without a condom in the past 12 months			Anal intercourse without a condom with last sex partner of discordant or unknown HIV status		
		n	(%)	<i>P</i> value <sup>a</sup>	n	(%)	<i>P</i> value <sup>a</sup>
<b>HIV-positive overall</b>	1113	804	(72.24)	<.001 <sup>b</sup>	478	(42.95)	<.001 <sup>b</sup>
<b>Race/Ethnicity</b>							
Black, non-Hispanic	63	46	(73.02)	.580	21	(33.33)	.009
Hispanic	162	110	(67.90)	.032	74	(45.68)	.780
White, non-Hispanic	810	590	(72.84)	REF	343	(42.35)	REF
Other or multiple races	78	58	(74.36)	.335	40	(51.28)	.027
<b>Age (years)</b>							
18-24	90	78	(86.67)	.060	56	(62.22)	.033
25-29	124	104	(83.87)	.085	66	(53.23)	.504
30-39	201	148	(73.63)	.041	94	(46.77)	.244
40 or older	698	474	(67.91)	REF	262	(37.54)	REF
<b>NHBS city resident<sup>c</sup></b>							
Yes	311	232	(74.60)	.016	119	(38.26)	.793
No	802	572	(71.32)	REF	359	(44.76)	REF
<b>Recruitment website type</b>							
Gay social networking	50	35	(70.00)	.987	25	(50.00)	.122
General gay interest	155	103	(66.45)	.139	57	(36.77)	.108
General social networking	439	291	(66.29)	REF	139	(31.66)	REF
Geospatial social networking	469	375	(79.96)	.001	257	(54.80)	.002
<b>HIV-negative or unknown overall</b>	9264	5673	(61.24)	REF	1262	(13.62)	REF
<b>Race/Ethnicity</b>							
Black, non-Hispanic	291	179	(61.51)	.297	55	(18.90)	.578
Hispanic	922	592	(64.21)	.934	183	(19.85)	.125
White, non-Hispanic	7266	4408	(60.67)	REF	901	(12.40)	REF
Other or multiple races	785	494	(62.93)	.651	123	(15.67)	.998
<b>Age (years)</b>							
18-24	1892	1217	(64.32)	.813	323	(17.07)	.001
25-29	1391	937	(67.36)	.019	193	(13.87)	.157
30-39	1717	1218	(70.94)	<.001	241	(14.04)	.613
40 or older	4264	2301	(53.96)	REF	505	(11.84)	REF
<b>NHBS city resident<sup>c</sup></b>							
Yes	2957	1784	(60.33)	.596	370	(12.51)	.737
No	6307	3889	(61.66)	REF	892	(14.14)	REF
<b>Recruitment website type</b>							
Gay social networking	752	365	(48.54)	<.001	95	(12.63)	.275
General gay interest	1803	1113	(61.73)	.134	205	(11.37)	.002
General social networking	4897	2926	(59.75)	REF	533	(10.88)	REF
Geospatial social networking	1812	1269	(70.03)	<.001	429	(23.68)	<.001

<sup>a</sup> Wald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared to a



referent (REF) group.

<sup>b</sup> Wald chi-square from multivariable logistic regression comparing behavior (yes versus no) among HIV-positive participants compared to HIV-negative or unknown serostatus participants. Model controlled for race/ethnicity, age, NHBS residency, and website type.

<sup>c</sup> NHBS = National HIV Behavioral Surveillance System

## Substance Use Behaviors

Illicit substance use in the past 12 months was more likely to be reported by HIV-positive participants than HIV-negative/unknown status participants (39.17% versus 26.85%, respectively;  $P < .001$ ; Table 4). Approximately half of participants reported binge drinking alcohol in the past 12 months, and there was no difference by participants' HIV status (55.53% for HIV-positive and 58.27% for HIV-negative/unknown;  $P = .681$ ).

Among participants who were HIV-positive, those 25-29 years of age were more likely to report using illicit drugs and binge drank alcohol compared with those  $\geq 40$  years of age (Table 4). More than half of those 25-29 years of age reported using illicit substances and more than three-quarters reported binge drinking alcohol in the past 12 months. Compared to participants recruited from a general social networking website, those from

gay general interest websites were less likely to report binge drank alcohol.

Among participants who were HIV-negative or unknown status, Hispanic participants were more likely and black or other/multiracial participants were less likely than white participants to report binge drinking alcohol (Table 4). Compared to participants  $\geq 40$  years of age, those 18-29 were more likely to report using illicit substances and binge drinking alcohol. Approximately one-third of these younger participants reported using illicit substances and three-quarters reported binge drinking alcohol in the past 12 months. Participants who resided in NHBS cities were also more likely to report using illicit substances and binge drinking. Compared to participants from the general social networking website, those from gay social networking websites were less likely to report substance use and those from a geospatial social networking website were more likely to report substance use.

**Table 4.** Substance using behaviors of MSM participants in the American Men's Internet Survey, United States, 2013.

Participant characteristics	n in sample	Substance use behaviors in the past 12 months					
		Used illicit drug			Binge drank alcohol		
		n	(%)	<i>P</i> value <sup>a</sup>	n	(%)	<i>P</i> value <sup>a</sup>
<b>HIV-positive overall</b>	1113	436	(39.17)	<.001 <sup>b</sup>	618	(55.53)	.681 <sup>b</sup>
<b>Race/Ethnicity</b>							
Black, non-Hispanic	63	27	(42.86)	.722	42	(66.67)	.434
Hispanic	162	68	(41.98)	.529	102	(62.96)	.981
White, non-Hispanic	810	304	(37.53)	REF	430	(53.09)	REF
Other or multiple races	78	37	(47.44)	.210	44	(56.41)	.435
<b>Age (years)</b>							
18-24	90	37	(41.11)	.360	66	(73.33)	.195
25-29	124	67	(54.03)	.007	100	(80.65)	<.001
30-39	201	94	(46.77)	.404	132	(65.67)	.502
40 or older	698	238	(34.10)	REF	320	(45.85)	REF
<b>NHBS city resident</b>							
Yes	311	131	(42.12)	.050	163	(52.41)	.625
No	802	305	(38.03)	REF	455	(56.73)	REF
<b>Recruitment website type</b>							
Gay social networking	50	15	(30.00)	.309	18	(36.00)	.025
General gay interest	155	57	(36.77)	.809	78	(50.32)	.271
General social networking	439	159	(36.22)	REF	227	(51.71)	REF
Geospatial social networking	469	205	(43.71)	.121	295	(62.90)	.131
<b>HIV-negative or unknown overall</b>	9264	2487	(26.85)	REF	5398	(58.27)	REF
<b>Race/Ethnicity</b>							
Black, non-Hispanic	291	75	(25.77)	.075	165	(56.70)	.007
Hispanic	922	275	(29.83)	.960	612	(66.38)	.007
White, non-Hispanic	7266	1912	(26.31)	REF	4187	(57.62)	REF
Other or multiple races	785	225	(28.66)	.336	434	(55.29)	.018
<b>Age (years)</b>							
18-24	1892	686	(36.26)	<.001	1349	(71.30)	<.001
25-29	1391	452	(32.49)	.006	1069	(76.85)	<.001
30-39	1717	521	(30.34)	.466	1143	(66.57)	.326
40 or older	4264	828	(19.42)	REF	1837	(43.08)	REF
<b>NHBS<sup>c</sup> city resident</b>							
Yes	2957	828	(28.00)	.002	1788	(60.47)	<.001
No	6307	1659	(26.30)	REF	3610	(57.24)	REF
<b>Recruitment website type</b>							
Gay social networking	752	118	(15.69)	<.001	339	(45.08)	<.001
General gay interest	1803	474	(26.29)	.176	1033	(57.29)	.773
General social networking	4897	1255	(25.63)	REF	2773	(56.63)	REF
Geospatial social networking	1812	640	(35.32)	<.001	1253	(69.15)	<.001

<sup>a</sup> Wald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared to a

referent (REF) group.

<sup>b</sup> Wald chi-square from multivariable logistic regression comparing behavior (yes versus no) among HIV-positive participants compared to HIV-negative or unknown serostatus participants. Model controlled for race/ethnicity, age, NHBS residency, and website type.

<sup>c</sup> NHBS = National HIV Behavioral Surveillance System

## HIV and STI Testing Behaviors

HIV testing behaviors were only examined among those who did not report being HIV-positive. Most of those participants (84.05%) had been previously tested for HIV infection, but less than half (44.20%) reported being tested in the past 12 months (Table 5). Compared to white participants, black participants were more likely to report ever having been tested. Compared to participants  $\geq 40$  years of age, those 18-24 years were less likely to report ever having been tested or having been tested in the past 12 months. Those 30-39 years were more likely to have been tested ever or in the past 12 months. Compared to participants recruited from the general social networking website, those from other websites had significant differences in reported HIV testing behaviors. Participants from gay social networking websites were less likely to report having been tested ever or in the past 12 months. Participants from general

gay interest websites and from a geospatial social networking website were more likely to report having been tested ever or in the past 12 months.

Among participants who were HIV-positive and got the randomized STI testing and diagnosis questions, 56.54% (216/382) had an STI test in the past 12 months and 19.89% (76/382) had any STI diagnosis: 9.16% (35/382) were diagnosed with gonorrhea, 7.07% (27/382) with Chlamydia and 9.69% (37/382) with syphilis. Among participants who were HIV-negative or unknown status, 24.48% (758/3096) had an STI test in the past 12 months and 4.98% (154/3096) had any STI diagnosis: 2.68% (83/3096) were diagnosed with gonorrhea, 2.62% (81/3096) with Chlamydia and 1.26% (39/3096) with syphilis. Compared to participants who were HIV-negative or of unknown status, those who were HIV-positive were significantly more likely to have been tested for and to have had any diagnosis of an STI (both  $P<.001$ ).

**Table 5.** HIV testing behaviors of HIV-negative or unknown status MSM participants in the American Men's Internet Survey, United States, 2013.

Participant characteristics	n in sample	Testing behaviors					
		HIV tested ever			HIV tested past 12 months		
		n	(%)	<i>P</i> value <sup>a</sup>	n	(%)	<i>P</i> value <sup>a</sup>
<b>Race/Ethnicity</b>							
Black, non-Hispanic	291	256	(87.97)	.009	160	(54.98)	.133
Hispanic	922	763	(82.75)	.541	458	(49.67)	.285
White, non-Hispanic	7266	6117	(84.19)	REF	3073	(42.29)	REF
Other or multiple races	785	649	(82.68)	.146	374	(47.64)	.786
<b>Age (years)</b>							
18-24	1892	1224	(64.69)	<.001	815	(43.08)	<.001
25-29	1391	1176	(84.54)	.295	688	(49.46)	.076
30-39	1717	1571	(91.50)	<.001	878	(51.14)	<.001
40 or older	4264	3814	(89.45)	REF	1684	(39.49)	REF
<b>NHBS <sup>b</sup> city resident</b>							
Yes	2957	2599	(87.89)	<.001	1403	(47.45)	<.001
No	6307	5186	(82.23)	REF	2662	(42.21)	REF
<b>Recruitment website type</b>							
Gay social networking	752	542	(72.07)	<.001	236	(31.38)	<.001
General gay interest	1803	1570	(87.08)	.008	730	(40.49)	<.001
General social networking	4897	4084	(83.40)	REF	1933	(39.47)	REF
Geospatial social networking	1812	1589	(87.69)	<.001	1166	(64.35)	<.001
Total	9264	7786	(84.05)		4095	(44.20)	

<sup>a</sup> Wald chi-square from multivariable logistic regression comparing behavior (yes versus no) among group with some characteristic compared to a referent (REF) group.

<sup>b</sup> NHBS = National HIV Behavioral Surveillance System

## Discussion

### Principal Findings

The first round of data collection for AMIS was successfully implemented and resulted in more than 10,000 surveys from a diverse sample of Internet using MSM residing in all US states. There were notable differences in key behavioral indicators sorted by recruitment website type. In particular, the geospatial social networking website produced a sample made up of participants with significantly different demographic characteristics and self-reported HIV status. Participants recruited from that site were also substantially more risky but also more likely to have been HIV tested. Future samples for AMIS and other analyses with this data will have to take these differences into consideration in study and analysis design.

One purpose of AMIS was to generate useful annual behavioral data to compare to NHBS-MSM which is only conducted every 3 years [3]. Understanding the differences between MSM recruited in the “in-person” NHBS surveys and our Internet-recruited surveys will allow correlation of NHBS and AMIS results and evaluation of trends in years between NHBS surveys. Compared to the most recent NHBS-MSM data from 2011, our study found a higher prevalence of all of the assessed risk behaviors, including for our sub-sample that lived in NHBS cities. Data from our study do not explain this difference, because where the few significant differences exist, AMIS participants that lived in NHBS cities were more likely than those that lived elsewhere to report risky behaviors. This risk difference between the two samples could be partially explained by differences between the demographic composition of the AMIS and NHBS samples, where NHBS had more participants who were younger and black. Our own study has shown that older participants were less likely to report risky behavior and other research has shown black MSM tend to report less unprotected sex and drug use [39]. Demographic differences between online and real-world samples have also been previously reported even where the geographic areas are the same, though unlike our study, most have found Internet surveys to have a higher proportion of younger participants than their comparison surveys [15,18,40]. The demographic characteristics of the AMIS sample and the WHBS pilot sample were more similar to one another than to NHBS also indicating that these differences in behaviors may be due less to geographic differences and more to sampling approach [35].

Regardless of those differences, our findings emphasize how annual AMIS data may complement those from the 3-year NHBS samples, providing timely and useful information for prevention program planning for MSM in many US states. In addition, the complementary data from AMIS may come at considerably reduced cost compared to the venue-recruited NHBS sample. The entire AMIS data collection and analysis support costs approximately \$15 per survey whereas NHBS costs at least \$1000 per survey based only on federal funding to local jurisdictions and not including CDC management or analysis costs [41]. NHBS is rightfully more expensive than AMIS because it involves a more comprehensive and detailed survey approach that requires full survey teams in each city to

conduct the in-person method, participant incentives, and laboratory expenses for rapid HIV testing.

Substantial proportions of AMIS participants reported sexual behaviors that may potentially pose a risk of HIV transmission. Over 40% of HIV-positive participants had anal sex without a condom with a potentially serodiscordant male partner. More than half of HIV-negative/unknown status participants also had anal sex without a condom, though most reported that they perceived their partners to be HIV-negative. Serostatus discussions between sex partners are an important part of HIV prevention, but previous studies have shown that those discussions may be based on inaccurate information because of high rates of undiagnosed HIV infection among MSM [39,42-44]. HIV-positive persons who are taking antiretroviral medications and have their HIV virus suppressed are also significantly less likely to transmit HIV to their sex partners, but we do not have this information for our participants [7]. Younger MSM were also significantly more likely to have had anal sex without a condom, a pattern also seen in the NHBS data [3]. For the youngest group in our study, 18-24 years of age, this is combined with a significantly increased likelihood that they are substance users and a reduced likelihood that they have ever been HIV tested. This presents a potentially heightened risk for HIV transmission in this group.

Compared to AMIS, NHBS had similar proportions who reported anal sex without a condom, but NHBS reported a substantially lower proportion of HIV-positive participants who had serodiscordant anal sex without a condom than our study (13% versus 43%, respectively) [3]. This may be explained by other differences in the samples/procedures (eg, demographics, self- versus interviewer-administered survey) or may be due to some fundamental difference in how sexual encounters are negotiated by HIV-positive MSM who were recruited from the internet. This indicator did not substantially differ between the HIV-negative/unknown status participants in the two studies. Similar to our findings, the EMIS study also found a significantly higher risk of non-concordant unprotected anal intercourse among HIV-positive participants compared to HIV-negative participants, but reported a substantially higher proportion of their participants overall had engaged in this behavior compared to our study (30% versus 17%, respectively) [34]. The timeframes for this behavior between our study and EMIS were not the same and may explain at least part of this difference; we examined behavior with last male sex partner and EMIS examined behavior with any male sex partner in the past 12 months. Collaborations should be explored to allow comparisons of non-concordant anal intercourse without condoms between these studies.

### Limitations

Several limitations to the AMIS methods and these analyses should be noted. First, AMIS data are not generalizable to all MSM in the US or to all MSM online. AMIS used a convenience sampling approach online and we cannot determine the degree or direction of response bias. Though we included several different types of websites to increase sample diversity, the websites still represent a small fraction of those that MSM likely use. Second, there was under-representation of black or African

American MSM in the AMIS sample, a problem common to Internet research [23]. This group is disproportionately impacted by HIV infection and the small AMIS sample size limits our ability to do more detailed analyses with these data. Third, the survey only involved self-report of behaviors. Though anonymous self-administered surveys such as AMIS may be less prone to obsequiousness bias [45-47], it is possible that less socially desirable responses may be under-reported (eg, anal sex without a condom) and more socially desirable responses may be over-reported (eg, recent HIV testing). Finally, the analyses presented here were only preliminary to illustrate the success of the AMIS method in generating key behavioral indicators. Although we report statistical tests in our behavioral analyses that controlled for some demographic characteristics, there were relatively few factors in the model which may not have resolved all confounding. Therefore interpretation regarding the independence of statistical relationships should

be made with caution until more detailed modeling can be conducted and reported.

### Future Directions

We are nearing completion of our second round of data collection of 10,000 surveys and intend to conduct the third round in the summer of 2015. The data we have collected to-date have been shared with state health departments in standardized reports to enable better planning for public health interventions (see [Multimedia Appendix 4](#) for an example report). We have also made individual state AMIS datasets available to each state's public health authorities so they can conduct further analyses for their own MSM residents. We have developed and deployed what will eventually become the largest ongoing Internet survey of MSM in the United States, and we envision that AMIS will become a useful tool in our joint endeavors to improve the health and wellbeing of this population.

### Acknowledgments

The study was funded by a grant from the MAC AIDS Fund and by the National Institutes of Health [P30AI050409] – the Emory Center for AIDS Research.

### Conflicts of Interest

Authors Sanchez and Sullivan are members of the Editorial Board of JMIR Public Health and Surveillance. However, they had no involvement in the editorial decision for this manuscript. It was reviewed and handled by an independent editor.

### Multimedia Appendix 1

AMIS 2013 questionnaire.

[\[PDF File \(Adobe PDF File\), 214KB-Multimedia Appendix 1\]](#)

### Multimedia Appendix 2

AMIS deduplication algorithm.

[\[PPTX File, 102KB-Multimedia Appendix 2\]](#)

### Multimedia Appendix 3

Characteristics of MSM participants in the American Men's Internet Survey by randomization to different question subsets, United States, 2013.

[\[PDF File \(Adobe PDF File\), 48KB-Multimedia Appendix 3\]](#)

### Multimedia Appendix 4

AMIS 2013 sample state report.

[\[PDF File \(Adobe PDF File\), 226KB-Multimedia Appendix 4\]](#)

### References

1. Centers for Disease Control and Prevention. 2014. Diagnoses of HIV Infection in the United States and Dependent Areas, 2012 URL: [http://www.cdc.gov/hiv/library/reports/surveillance/2012/surveillance\\_Report\\_vol\\_24.html](http://www.cdc.gov/hiv/library/reports/surveillance/2012/surveillance_Report_vol_24.html) [accessed 2015-02-03] [[WebCite Cache ID 6W4bEeOFn](#)]
2. Ward BW, Dahlhamer JM, Galinsky AM, Joestl SS. National Health Statistics Reports.: Centers for Disease Control and Prevention; 2014 Jul 15. Sexual orientation and health among US adults: national health interview survey, 2013 URL: <http://www.cdc.gov/nchs/data/nhsr/nhsr077.pdf> [accessed 2015-04-13] [[WebCite Cache ID 6Xm0UMOXp](#)]
3. Centers for Disease Control and Prevention. HIV testing and risk behaviors among gay, bisexual, and other men who have sex with men - United States. Morbidity and Mortality Weekly Report 2013;62(47):958-962.



4. Johnson AS, Hall HI, Hu X, Lansky A, Holtgrave DR, Mermin J. Trends in diagnoses of HIV infection in the United States, 2002-2011. *JAMA* 2014;312(4):432-434. [doi: [10.1001/jama.2014.8534](https://doi.org/10.1001/jama.2014.8534)] [Medline: [25038362](#)]
5. Centers for Disease Control and Prevention. 2012. CDC Fact Sheet: New HIV Infections in the United States URL: <http://www.cdc.gov/nchhstp/newsroom/docs/2012/HIV-Infections-2007-2010.pdf> [WebCite Cache ID 6VsLm07w]
6. Grant RM, Lama JR, Anderson PL, McMahan V, Liu AY, Vargas L, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med* 2010 Dec 30;363(27):2587-2599 [FREE Full text] [doi: [10.1056/NEJMoa1011205](https://doi.org/10.1056/NEJMoa1011205)] [Medline: [21091279](#)]
7. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N, Pilotto Jose H S, HPTN 052 Study Team. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med* 2011 Aug 11;365(6):493-505 [FREE Full text] [doi: [10.1056/NEJMoa1105243](https://doi.org/10.1056/NEJMoa1105243)] [Medline: [21767103](#)]
8. Sullivan PS, Carballo-Diéguez A, Coates T, Goodreau SM, McGowan I, Sanders EJ, et al. Successes and challenges of HIV prevention in men who have sex with men. *Lancet* 2012 Jul 28;380(9839):388-399 [FREE Full text] [doi: [10.1016/S0140-6736\(12\)60955-6](https://doi.org/10.1016/S0140-6736(12)60955-6)] [Medline: [22819659](#)]
9. Centers for Disease Control and Prevention. 2014. Laboratory testing for the diagnosis of HIV infection : updated recommendations URL: <http://www.cdc.gov/hiv/pdf/HIVtestingAlgorithmRecommendation-Final.pdf> [accessed 2015-02-03] [WebCite Cache ID 6W4bW0D9q]
10. Centers for Disease Control and Prevention. 2014. Rapid HIV tests suitable for use in non-clinical settings URL: [http://www.cdc.gov/hiv/pdf/testing\\_nonclinical\\_clia-waived-tests.pdf](http://www.cdc.gov/hiv/pdf/testing_nonclinical_clia-waived-tests.pdf) [accessed 2015-02-03] [WebCite Cache ID 6W4bZao2M]
11. Pew Research Internet Project. Social Media Update 2014 URL: <http://www.pewinternet.org/2015/01/09/social-media-update-2014/> [accessed 2015-01-20] [WebCite Cache ID 6VjXjUHqX]
12. Bolding G, Davis M, Hart G, Sherr L, Elford J. Where young MSM meet their first sexual partner: the role of the Internet. *AIDS Behav* 2007 Jul;11(4):522-526. [doi: [10.1007/s10461-007-9224-9](https://doi.org/10.1007/s10461-007-9224-9)] [Medline: [17347876](#)]
13. Grosskopf NA, Harris JK, Wallace BC, Nanin JE. Online sex-seeking behaviors of men who have sex with men in New York City. *Am J Mens Health* 2011 Sep;5(5):378-385. [doi: [10.1177/1557988310372801](https://doi.org/10.1177/1557988310372801)] [Medline: [20798150](#)]
14. Horvath KJ, Rosser B R Simon, Remafedi G. Sexual risk taking among young internet-using men who have sex with men. *Am J Public Health* 2008 Jun;98(6):1059-1067. [doi: [10.2105/AJPH.2007.111070](https://doi.org/10.2105/AJPH.2007.111070)] [Medline: [18445804](#)]
15. Sanchez T, Smith A, Denson D, DiNenno E, Lansky A. Internet-based methods may reach higher risk men who have sex with men not reached through venue-based sampling. *Open AIDS* 2012;6(Supplement 1):99-105.
16. Rhodes SD, DiClemente RJ, Cecil H, Hergenrather KC, Yee LJ. Risk among men who have sex with men in the United States: a comparison of an Internet sample and a conventional outreach sample. *AIDS Educ Prev* 2002 Feb;14(1):41-50. [Medline: [11900109](#)]
17. Bull SS, Lloyd L, Rietmeijer C, McFarlane M. Recruitment and retention of an online sample for an HIV prevention intervention targeting men who have sex with men: the Smart Sex Quest Project. *AIDS Care* 2004 Nov;16(8):931-943. [doi: [10.1080/09540120412331292507](https://doi.org/10.1080/09540120412331292507)] [Medline: [15511725](#)]
18. Elford J, Bolding G, Davis M, Sherr L, Hart G. Web-based behavioral surveillance among men who have sex with men: a comparison of online and offline samples in London, UK. *J Acquir Immune Defic Syndr* 2004 Apr 1;35(4):421-426. [Medline: [15097159](#)]
19. Chiasson MA, Hirshfield S, Remien RH, Humberstone M, Wong T, Wolitski RJ. A comparison of on-line and off-line sexual risk in men who have sex with men: an event-based on-line survey. *J Acquir Immune Defic Syndr* 2007 Feb 1;44(2):235-243. [doi: [10.1097/QAI.0b013e31802e298c](https://doi.org/10.1097/QAI.0b013e31802e298c)] [Medline: [17179769](#)]
20. Berg RC. Barebacking among MSM Internet users. *AIDS Behav* 2008 Sep;12(5):822-833. [doi: [10.1007/s10461-007-9281-0](https://doi.org/10.1007/s10461-007-9281-0)] [Medline: [17676278](#)]
21. Rosser B R Simon, Miner MH, Bockting WO, Ross MW, Konstan J, Gurak L, et al. HIV risk and the internet: results of the Men's INternet Sex (MINTS) Study. *AIDS Behav* 2009 Aug;13(4):746-756 [FREE Full text] [doi: [10.1007/s10461-008-9399-8](https://doi.org/10.1007/s10461-008-9399-8)] [Medline: [18512143](#)]
22. Gilbert LK, Levandowski BA, Scanlon KE, Peterson RS. A comparison of hepatitis A and hepatitis B measures among vaccinated and susceptible online men who have sex with men. *Int J STD AIDS* 2010 Jun;21(6):400-405. [doi: [10.1258/ijisa.2009.009010](https://doi.org/10.1258/ijisa.2009.009010)] [Medline: [20606219](#)]
23. Sullivan PS, Khosropour CM, Luisi N, Amsden M, Coggia T, Wingood GM. Bias in online recruitment and retention of racial and ethnic minority men who have sex with men. *Journal of Medical Internet Research* 2011;13(2).
24. Krakower DS, Mimiaga MJ, Rosenberger JG, Novak DS, Mitty JA, White JM. Limited Awareness and Low Immediate Uptake of Pre-Exposure Prophylaxis among Men Who Have Sex with Men Using an Internet Social Networking Site. *PloS One* 2012;7(3).
25. Rosenberger JG, Reece M, Schick V, Herbenick D, Novak DS, Van Der Pol Barbara, et al. Condom use during most recent anal intercourse event among a U.S. sample of men who have sex with men. *J Sex Med* 2012 Apr;9(4):1037-1047. [doi: [10.1111/j.1743-6109.2012.02650.x](https://doi.org/10.1111/j.1743-6109.2012.02650.x)] [Medline: [22353190](#)]
26. Berg RC, Tikkanen R, Ross MW. Barebacking among men who have sex with men recruited through a Swedish website: associations with sexual activities at last sexual encounter. *European Communicable Disease Bulletin* 2013;18(13):06.

27. Greacen T, Friboulet D, Blachier A, Fugon L, Hefez S, Lorente N, et al. Internet-using men who have sex with men would be interested in accessing authorised HIV self-tests available for purchase online. *AIDS Care* 2013;25(1):49-54. [doi: [10.1080/09540121.2012.687823](https://doi.org/10.1080/09540121.2012.687823)] [Medline: [22670681](#)]
28. Yu G, Wall MM, Chiasson MA, Hirshfield S. Complex drug use patterns and associated HIV transmission risk behaviors in an Internet sample of U.S. men who have sex with men. *Arch Sex Behav* 2015 Feb;44(2):421-428. [doi: [10.1007/s10508-014-0337-8](https://doi.org/10.1007/s10508-014-0337-8)] [Medline: [25104104](#)]
29. Dasgupta S, Vaughan AS, Kramer MR, Sanchez TH, Sullivan PS. Use of a Google Map Tool Embedded in an Internet Survey Instrument: Is it a Valid and Reliable Alternative to Geocoded Address Data? *JMIR Res Protoc* 2014;3(2):e24 [FREE Full text] [doi: [10.2196/resprot.2946](https://doi.org/10.2196/resprot.2946)] [Medline: [24726954](#)]
30. Sineath RC, Finneran C, Sullivan P, Sanchez T, Smith DK, Griensven FV, et al. Knowledge of and interest in using preexposure prophylaxis for HIV prevention among men who have sex with men in Thailand. *J Int Assoc Provid AIDS Care* 2013;12(4):227-231. [doi: [10.1177/2325957413488184](https://doi.org/10.1177/2325957413488184)] [Medline: [23708677](#)]
31. Santos GM, Do T, Beck J, Makofane K, Arreola S, Pyun T, et al. Syndemic conditions associated with increased HIV risk in a global sample of men who have sex with men. *Sex Transm Infect* 2014 May;90(3):250-253. [doi: [10.1136/sextrans-2013-051318](https://doi.org/10.1136/sextrans-2013-051318)] [Medline: [24431183](#)]
32. Hickson F, Bonell C, Hargreaves J, Reid D, Weatherburn P. HIV Testing and HIV Serostatus-Specific Sexual Risk Behaviour Among Men Who Have Sex with Men Living in England and Recruited Through the Internet in 2001 and 2008. *Sex Res Soc Policy* 2012 Dec 2;10(1):15-23. [doi: [10.1007/s13178-012-0106-1](https://doi.org/10.1007/s13178-012-0106-1)]
33. Sigma Research. The Gay Men's Sex Survey. 2015. Sample Sizes URL: <http://www.sigmaresearch.org.uk/gmss/survey/sample> [accessed 2015-01-27] [WebCite Cache ID 6VsiSRKo6]
34. The EMISN. Findings from 38 countries. EMIS 2010: The European Men-Who-Have-Sex-With-Men Internet Survey URL: [http://www.emis-project.eu/sites/default/files/public/publications/emis-2010\\_european\\_msm\\_internet\\_survey\\_38\\_countries\\_v5.pdf](http://www.emis-project.eu/sites/default/files/public/publications/emis-2010_european_msm_internet_survey_38_countries_v5.pdf) [accessed 2015-01-21] [WebCite Cache ID 6VjZO8w7m]
35. Sanchez T, Smith A, Denson D, Dinunno E, Lansky A. Developing a Web-Based HIV Behavioral Surveillance Pilot Project Among Men Who Have Sex with Men. *Open AIDS J* 2012;6:224-231 [FREE Full text] [doi: [10.2174/1874613601206010224](https://doi.org/10.2174/1874613601206010224)] [Medline: [23091580](#)]
36. Centers for Disease Control and Prevention. 2011. National HIV Behavioral Surveillance System Round 3 Questionnaire URL: [http://www.cdc.gov/hiv/pdf/NHBS\\_Round3\\_OMBApprovedQuestionnaire-508remediated.pdf](http://www.cdc.gov/hiv/pdf/NHBS_Round3_OMBApprovedQuestionnaire-508remediated.pdf) [accessed 2015-01-21] [WebCite Cache ID 6VjZqrbD0]
37. Sanchez T, Finlayson T, Drake A, Behel S, Cribbin M, Dinunno E, Centers for Disease ControlPrevention (CDC). Human immunodeficiency virus (HIV) risk, prevention, and testing behaviors--United States, National HIV Behavioral Surveillance System: men who have sex with men, November 2003-April 2005. *MMWR Surveill Summ* 2006 Jul 7;55(6):1-16 [FREE Full text] [Medline: [16826162](#)]
38. National Institute of Alcohol Abuse and Alcoholism. 2004. NIAAA Council Approves Definition of Binge Drinking URL: [http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter\\_Number3.pdf](http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter_Number3.pdf) [accessed 2015-03-11] [WebCite Cache ID 6WxahUEAA]
39. Maulsby C, Millett G, Lindsey K, Kelley R, Johnson K, Montoya D, et al. HIV among Black men who have sex with men (MSM) in the United States: a review of the literature. *AIDS Behav* 2014 Jan;18(1):10-25. [doi: [10.1007/s10461-013-0476-2](https://doi.org/10.1007/s10461-013-0476-2)] [Medline: [23620241](#)]
40. Evans AR, Wiggins RD, Mercer CH, Bolding GJ, Elford J. Men who have sex with men in Great Britain: comparison of a self-selected internet sample with a national probability sample. *Sex Transm Infect* 2007 Jun;83(3):200-5; discussion 205 [FREE Full text] [doi: [10.1136/sti.2006.023283](https://doi.org/10.1136/sti.2006.023283)] [Medline: [17135330](#)]
41. Centers for Disease Control and Prevention. Division of HIV/AIDS Prevention Awards by State, 2012 URL: [http://www.cdc.gov/hiv/pdf/policies\\_funding\\_awards2012.pdf](http://www.cdc.gov/hiv/pdf/policies_funding_awards2012.pdf) [accessed 2015-03-11] [WebCite Cache ID 6WxF9BeJe]
42. Winter AK, Sullivan PS, Khosropour CM, Rosenberg ES. Discussion of HIV status by serostatus and partnership sexual risk among internet-using MSM in the United States. *J Acquir Immune Defic Syndr* 2012 Aug 15;60(5):525-529 [FREE Full text] [doi: [10.1097/QAI.0b013e318257d0ac](https://doi.org/10.1097/QAI.0b013e318257d0ac)] [Medline: [22549381](#)]
43. Sullivan PS, Wall KM, O'Hara B, Jones J, Barnes J, DiClemente R, et al. The prevalence of undiagnosed HIV serodiscordance among male couples presenting for HIV testing. *Arch Sex Behav* 2014 Jan;43(1):173-180 [FREE Full text] [doi: [10.1007/s10508-013-0214-x](https://doi.org/10.1007/s10508-013-0214-x)] [Medline: [24233391](#)]
44. Millett GA, Ding H, Marks G, Jeffries WL, Bingham T, Lauby J, et al. Mistaken assumptions and missed opportunities: correlates of undiagnosed HIV infection among black and Latino men who have sex with men. *J Acquir Immune Defic Syndr* 2011 Sep 1;58(1):64-71. [doi: [10.1097/QAI.0b013e31822542ad](https://doi.org/10.1097/QAI.0b013e31822542ad)] [Medline: [21654500](#)]
45. Langhaug LF, Sherr L, Cowan FM. How to improve the validity of sexual behaviour reporting: systematic review of questionnaire delivery modes in developing countries. *Trop Med Int Health* 2010 Mar;15(3):362-381 [FREE Full text] [doi: [10.1111/j.1365-3156.2009.02464.x](https://doi.org/10.1111/j.1365-3156.2009.02464.x)] [Medline: [20409291](#)]
46. Hewett PC, Mensch BS, Ribeiro Manoel Carlos S de A, Jones HE, Lippman SA, Montgomery MR, van de Wijgert Janneke H H M. Using sexually transmitted infection biomarkers to validate reporting of sexual behavior within a randomized,

- experimental evaluation of interviewing methods. Am J Epidemiol 2008 Jul 15;168(2):202-211 [[FREE Full text](#)] [doi: [10.1093/aje/kwn113](#)] [Medline: [18525081](#)]
47. Metzger DS, Koblin B, Turner C, Navaline H, Valenti F, Holte S, et al. Randomized controlled trial of audio computer-assisted self-interviewing: utility and acceptability in longitudinal studies. HIVNET Vaccine Preparedness Study Protocol Team. Am J Epidemiol 2000 Jul 15;152(2):99-106 [[FREE Full text](#)] [Medline: [10909945](#)]

## Abbreviations

**AMIS:** American Men's Internet Survey  
**CDC:** Centers for Disease Control and Prevention  
**EMIS:** European MSM Internet Survey  
**HIV:** human immunodeficiency virus  
**IP:** Internet protocol  
**MSM:** men who have sex with men  
**NHBS:** National HIV Behavioral Surveillance System  
**STI:** sexually transmitted infection  
**WHBS:** Web-based HIV Behavioral Surveillance System

*Edited by G Eysenbach; submitted 03.02.15; peer-reviewed by R Pendse, O Martinez; comments to author 25.02.15; revised version received 17.03.15; accepted 17.03.15; published 17.04.15*

*Please cite as:*

Sanchez TH, Sineath RC, Kahle EM, Tregear SJ, Sullivan PS

*The Annual American Men's Internet Survey of Behaviors of Men Who Have Sex With Men in the United States: Protocol and Key Indicators Report 2013*

*JMIR Public Health Surveill* 2015;1(1):e3

URL: <http://publichealth.jmir.org/2015/1/e3/>

doi: [10.2196/publichealth.4314](#)

PMID: [27227126](#)

©Travis Howard Sanchez, R Craig Sineath, Erin M Kahle, Stephen James Tregear, Patrick Sean Sullivan. Originally published in JMIR Public Health and Surveillance (<http://publichealth.jmir.org>), 17.04.2015. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Public Health and Surveillance, is properly cited. The complete bibliographic information, a link to the original publication on <http://publichealth.jmir.org>, as well as this copyright and license information must be included.